



# Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

---

## Monoclonal Antibody To Human MRP8 (S100A8) Calgranulin A - Marker For A Subpopulation Of Inflammatory Leukocytes

Monoclonal antibody 8-5C2 identifies MRP8 (also named S100A8 or Calgranulin A), the  $\text{Ca}^{2+}$ -binding light subunit of the inflammatory L-1 protein complex. MRP8 forms  $\text{Ca}^{2+}$  dependent dimers or complexes with MRP14 (S100A9, Calgranulin B). It also forms disulfide-linked homodimers under the influence of hypochlorite, a process thought to abrogate the chemotactic property of MRP8. The antibody is useful in various immunological techniques. Histological and serological data indicate that MRP8 is associated with chronic stages of inflammatory diseases.

---

<b>Product Number:</b>	T-1030
<b>Clone:</b>	8-5C2
<b>Host species, isotype:</b>	Mouse IgG1
<b>Quantity:</b>	100 $\mu$ g
<b>Format:</b>	Affinity purified, lyophilized
	Reconstitute by adding 0.5ml distilled water. This stock solution contains 0.2mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), 5mg/ml bovine serum albumin (BSA) as a stabilizer and 0.09% sodium azide as a preservative.
<b>Stability:</b>	Original vial: 1 year at 4° - 8°C Stock solution or aliquots thereof: 1 year at -20°C. Avoid repeated thawing and freezing.
<b>Applications:</b>	Tested for immunohistochemistry (IHC) and ELISA; has been described to work in dot blots, not in FACS. <b>Approximate working dilution for IHC:</b> Frozen sections: 1 - 2 $\mu$ g/ml (1:100 - 1:200) Paraffin sections: 5 - 10 $\mu$ g/ml (1:20 - 1:50); no pretreatment for antigen retrieval necessary; microwave treatment in 0.01M citrate pH 6.0 may enhance the reactivity. Optimal dilutions should be determined by the end user. Suggested positive control: Human tonsil.
<b>Immunogen:</b>	Cultured human monocytes.
<b>Antigen, epitope:</b>	The antigen is MRP8, the epitope is suspected in the central portion of the peptide.



# Peninsula Laboratories, LLC

**A Member of the Bachem Group**

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

**Antigen distribution:**

**Isolated cells:** The antigen is found in granulocytes and monocytes but not in other blood cells. In cultured monocytes, maximum MRP8 is expressed after 3 - 4 days. Myeloid leukaemia stain positive.

**Tissue sections:** MRP8 is found in a distinct subpopulation of inflammatory perivascular infiltrates of the myelo-monocytic lineage. Macrophages synthesise MRP8 increasingly during the late stages of inflammation. A low MRP8 (and high MRP-14) expression by macrophages was also reported in granulomatous diseases such as tuberculosis and sarcoidis. In non-granulomatous chronic inflammatory diseases such as chronic rheumatoid arthritis or chronic rejection after allograft transplantation, MRP8 and MRP14 positive cells consist of different subpopulations.

**Specificity:**

**Human:** MRP8 in stimulated monocytes and macrophages in late phase or chronic inflammation.

**Other:** The antibody does not react with rat tissues.

---

## Selected references

Odink, K., et al.: Two calcium-binding proteins in infiltrate macrophages of rheumatoid arthritis. *Nature* **330**, 80 (1987).

Brandtzaeg, P et al.: Mac387 antibody and detection of formalin resistant myelomonocytic L1 protein. *J.Clin.Pathol.*: **41**, 963- (1988).

Zwadlo, G.,et al.: Two calcium-binding proteins associated with specific stages of myeloid cell differentiation are expressed by subsets of macrophages in inflammatory tissues. *Clin.Exp.Immunol.*: **72**, 510- (1988).

Murao, S. et al.: A Protein Containing the Cystic Fibrosis Antigen is an Inhibitor of Protein Kinases. *J. Biological Chem.*: **254**, (14) 8356 - 8360 (1989).

Mues, B., et al.: Phenotyping of macrophages with monoclonal antibodies in endomyocardial biopsies as a new approach to diagnosis of myocarditis. *Eur. Heart Journal*: **11**, 619- (1990).

Delabie, J., et al.: Differential expression of the calcium-binding proteins MRP8 and MRP14 in granulomatous conditions: an immunohistochemical study. *Clin.Exp.Immunol.*: **81**, 123- (1990).

Teigelkamp, S., et al.: Calcium dependent Complex Assembly of the Myeloid Differentiation Proteins MRP-8 and MRP-14. *J. Biological Chem.*: **266**, (20), 13462 - 13467, (1991).

Sorg, C.: *Macrophages in Inflammation*. Regensberg & Biermann. ISBN 3-924469-23-7: 23-35 (1988).

For in vitro research only. Caution: this product contains sodium azide, a poisonous and hazardous substance.