



## Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

---

### Monoclonal Antibody to Human MCA

#### Mucin-Like Carcinoma Antigen, Marker For Mucin Producing Cells

Monoclonal antibody b-12 is useful for identifying mucin-like carcinoma antigen (MCA) produced by various tumours and certain healthy glandular cells. In combination with other markers for inflammation staging or investigating neo-vascularization processes, b-12 is a valuable tool for studying tumour growth or regression. MCA is a 350kDa glycoprotein with the typical biochemical characteristics of mucin-like glycoproteins (sialomucins) which protect surfaces. Antibody b-12 binds to the protein backbone of MCA, not to the large number of carbohydrate side chains.

---

<b>Product Number:</b>	T-1301
<b>Clone:</b>	b-12
<b>Host species, isotype:</b>	Mouse IgG1
<b>Quantity:</b>	50µg
<b>Format:</b>	Affinity purified, lyophilized
	Reconstitute by adding 0.5ml distilled water. This stock solution contains 0.1mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), 2mg/ml bovine serum albumin (BSA) as a stabilizer and 0.05% sodium azide as a preservative.
<b>Stability:</b>	Original vial: 1 year at 4° - 8°C Stock solution or aliquots thereof: 1 year at -20°C. Avoid repeated thawing and freezing.
<b>Applications:</b>	Tested for immunohistochemistry (IHC). <b>Approximate working dilution for IHC:</b> Frozen sections: 0.5µg/ml (1:200) Paraffin sections: 4µg/ml (1:25); Proteinase K pretreatment for antigen retrieval recommended. Optimal dilutions should be determined by the end user. Suggested positive control: Human uterus.
<b>Immunogen:</b>	Breast carcinoma cell lines.
<b>Antigen, epitope:</b>	MCA consists of a polymorphic family of glycoproteins. The b-12 related antigenic epitope is located in the more constant region of MCA.



# Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

**Antigen distribution:** In contrast to MCA producing tumours, the b-12 related antigen is only located at the MCA producing sites such as glandular cell surfaces or glandular tubuli. In MCA producing tumours, where cells become disorganized, the b-12 antigen is secreted into stromal tissue and blood vessels.

### ***b-12 Reaction Pattern on Human Tissues:***

Healthy Tissues		Cancerous Tissues	
Transitional epithelium	3 / 3	Breast	122 / 122
Kidney	13 / 13	Uterus: Endometrium	10 / 10
Fallopian tube	2 / 2	Cervix, squamous cells	2 / 2
Uterus	5 / 5	Ovary Mucinous	4 / 4
Prostate	6 / 9	Serous	2 / 2
Epididymis	4 / 4	Testis Malignant teratoma	7 / 7
Bronchus	13 / 13	Kidney Clear cell	15 / 15
Sebaceous and sweat glands	6 / 6	Lung Bronchiolo-alveolar	6 / 6
Salivary glands	3 / 4	Adenosquamous	2 / 2
Stomach	6 / 8	Stomach Adenocarcinoma	8 / 9
Breast	23 / 23	Colon Adenocarcinoma	24 / 36

(from Zenklusen et al. 1988)

**Specificity:** **Human:** MCA producing cells.

**Other species:** not tested.

---

### **Selected references**

Staehli, C. et al.: Monoclonal antibodies against antigens on breast cancer cells. *Experientia* **41**: 1377 (1985)

Zenklusen, H.R. et al.: The immunohistochemical reactivity of a new anti-epithelial antibody (mAb b-12) against breast carcinoma and other normal and neoplastic human tissues. *Virchows Arch A Pathol Anat* **413**: 3 (1988)

Maurer, A. & Burckhardt, J.: Biochemistry and molecular biology of MCA. *Int. J. Biol. Markers* **8**: 108-112 (1993).

For in vitro research only. Caution: this product contains sodium azide, a poisonous and hazardous substance.