



Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

Monoclonal Antibody To Mouse Monocytes / Macrophages Marker With Broad Specificity

Monoclonal antibody MOMA-2 is a useful marker for the broad detection of monocytes and macrophages in all mouse strains. In combination with the anti F4/80 marker BM8 (product T-2006) it allows a precise characterisation of tissue fixed macrophages in various organs. The antibody stains a mature macrophage subset, monocytes and a few precursors in bone marrow. Dendritic cells show low to intermediate expression. The staining shows close correlation with expression of acid phosphatase in tissue sections. MOMA-2 is predominantly expressed in the cytoplasm, but is also present on the cell surface.

Product Number:	T-2007
Clone:	MOMA-2
Host species, isotype:	Rat IgG2b
Quantity:	200µg
Format:	Affinity purified, lyophilized Reconstitute by adding 0.5ml distilled water. This stock solution contains 0.4mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), 5mg/ml bovine serum albumin (BSA) as a stabilizer and 0.1% sodium azide as a preservative.
Stability:	Original vial: 1 year at 4° - 8°C Stock solution or aliquots thereof: 1 year at -20°C. Avoid repeated thawing and freezing.
Applications:	Tested for immunohistochemistry (IHC); has been described to work in FACS (after permeabilisation). Approximate working dilution for IHC: Frozen sections: 0.5-1µg/ml (1:400 - 1:800) Paraffin sections: does not react on routinely processed paraffin sections. Optimal dilutions should be determined by the end user. Suggested positive control: Mouse spleen.
Immunogen:	Mouse lymph node stroma.
Antigen, epitope:	The antigen is a (glyco-)protein of 140kDa m.w. which is located within the cytoplasm and on the cell surface. The epitope has not been further characterized.



Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

Antigen distribution:

Isolated cells: In the cytospin preparation of thioglycollate stimulated peritoneal exudate cells MOMA-2 detects an antigen as distinct cytoplasmic spots. MOMA-2 detects monocytes of the peripheral blood and a subpopulation of bone marrow cells

Tissue sections MOMA-2 detects typical tissue macrophages as does the anti F4/80-specific clone BM8 (pan macrophage marker). However, different staining patterns are visible as shown below. The most predominant difference can be observed in T-cell areas and follicles of peripheral lymphoid organs where the anti F4/80 clone BM8 is negative.

Comparison of different mature macrophage markers

Product number	MOMA-2 T-2007	BM8 (anti F4/80) T-2006	ER-BMDM 1 T-2015
Monocytes	+	+	-
Kupffer cells	+	+	-
Langerhans cells	+/-	+	
Tingible body macrophages	+	-	
Interdigitating cells	+/-	-	+
Dendritic cells	+/-	-	+
Microglial cells	-		-
Marginal zone macrophages	-	-	
Marginal metallophilic cells	-	-	-
Pneumocytes type II			+
Alveolar lavage cells		66%	26%
Resident peritoneal cells (PCs)		51%	34%
Thioglycollate elicited PCs time after injection: 4hours		81%	79%
time after injection: 8 hours		28%	15%
Bone Marrow (BM) cells	14%	37%	5%
BM cells after 7 days with M-CSF	30%	96%	91%

Kraal et al. (1987) modified and P.J.M. Leenen personal communication

Specificity:

Mouse: monocytes and macrophages.

Other species: not tested.

Selected references

BREEL, M., MEBIUS, R.E., KRAAL, G.: Dendritic cells of the mouse recognized by two monoclonal antibodies. *Eur. J. Immunol.*: 17, 1555 - 1559 (1987).

KRAAL, G., REP, M., JANSE, M.: Macrophages in T and B Cell Compartments and Other Tissue Macrophages Recognized by Monoclonal Antibody MOMA-2; An Immunohistochemical Study. *Scand. J. Immunol.*: 26, 653 - 661 (1987).

For in vitro research only. Caution: this product contains sodium azide, a poisonous and hazardous substance.