



Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

FITC Labeled Monoclonal Antibody To Mouse CD205

Marker For Mouse Nonlymphoid Dendritic Cells, DEC-205

Monoclonal antibody NLDC-145 identifies Ia positive interdigitating cells, veiled cells and Langerhans cells of the skin and their *in vitro* counterparts. The antigen is expressed at high levels by dendritic cells and thymic epithelial cells. The antigen detected by NLDC-145 is an integral membrane glycoprotein with an apparent mass of 205kDa, also known as DEC-205. DEC-205 is apparently a receptor involved in antigen-processing by dendritic cells.

Product Number:	T-2023
Clone:	NLDC-145
Host species, isotype:	Rat IgG2a
Quantity:	200µg
Format:	Affinity purified, FITC labelled, liquid Supplied as 1ml solution. This stock solution contains 0.2mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), 10mg/ml bovine serum albumin (BSA) as a stabilizer and 0.09% sodium azide as a preservative.
Stability:	Original vial: 6 months at 4° - 8°C
Applications:	Has been described to work in FACS. Approximate working dilution: Optimal dilutions should be determined by the end user. Suggested positive control: GM-CSF and IL-4 stimulated bone marrow cells
Immunogen:	Mouse lymph node tissue.
Antigen, epitope:	The antigen is a protein of 205kDa molecular weight (DEC-205) which is localized on the cell surface and intracellularly. The epitope has not been further characterized.



Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

Antigen distribution:

Organ	NLDC-145 staining	Cell type and localization
Spleen	+	IDC in inner PALS
Lymph node	+	IDC in paracortex VC in subcapsular sinus
Peyer's patch	+	IDC in interfollicular T cell areas Villum epithelium, isolated cells in submucosa (VC)
Thymus	+	IDC in medulla Cortical epithelium
Skin	+	Langerhans cells
Brain, Kidney, Liver, Heart	-	
In vitro isolated cells		
Blood, bone marrow	-	
Peritoneal cells	-	
Peritoneal exudate cells*	+	some positive cells (VC?)

* Peritoneal exudate cells were harvested 4 days after intraperitoneal thioglycollate injection.

Tests were carried out on BALB/c and C₃D₂F₁ mouse strains.

(G.Kraal et al. see ref. 1, modified)

Specificity:

Mouse: nonlymphoid dendritic cells: interdigitating cells (IDC), veiled cells and Langerhans cells, thymic epithelial cells.

Other species: not tested.

Selected references

Kraal, G., M. Breel, M. Janse, G. Bruin: Langerhans' cells, veiled cells, and interdigitating cells in the mouse recognized by a monoclonal antibody. *J Exp Med* **163**: 981-987 (1986)

Breel, M., R.E. Mebius, G. Kraal: Dendritic cells of the mouse recognized by two monoclonal antibodies. *Eur J Immunol* **17**: 1555-1559 (1987).

Swiggard, W.J., A. Mirza, M.C. Nussenzweig, R.M. Steinman: DEC-205, a 205kDa protein abundant on mouse dendritic cells and thymic epithelium that is detected by the monoclonal antibody NLDC-145: purification, characterization, and N-terminal amino acid sequence. *Cellular Immunology* **165**, 302-311 (1995).

Martinez del Hoyo, G., P. Martín, H. Hernández Vargas, S. Ruiz, C. Fernández Arias, C. Ardavín: Characterization of a common precursor population for dendritic cells. *Nature* **415**: 1043-47 (2002)

For in vitro research only. Caution: this product contains sodium azide, a poisonous and hazardous substance.