

PENINSULA LABORATORIES, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070 Tel: (800) 922-1516 • (650) 592-5392 Fax: (650) 595-4071

www.bachem.com

Cat. No. T-4041

Rabbit anti-β-Endorphin (human)

Specifications:

Preparation: This polyclonal antibody was collected from rabbits immunized

with a synthetic peptide as the immunogen.

Amount: Each vial contains 50 ul of whole serum in lyophilized form.

Reconstitute with 50 ul of distilled water for the equivalent of

undiluted antiserum.

Storage: This antibody is supplied as a lyophilized powder. Store at 2° to

8°C upon receipt. For long-term storage, store lyophilized powder at -20°C. Once reconstituted, the materials should be stored frozen in aliquot at -20°C to -70°C. **Avoid repeated**

freeze/thaw cycles.

Applications: Immunohistochemistry, Immunocytochemistry, and other

applications.

Sample Types: The antibody can react with a variety of sample preparation

including

frozen tissue sections, paraffin embedded tissue sections and fixed

cell cultures.

Recommended starting dilution for Immunohistochemical staining:

Immunofluorescence: 1:250 to 1:16000

PAP or ABC: 1:250 to 1:16000 LAB-SA: 1:250 to 1:16000

It is advised for each laboratory to determine optimal dilutions for a specific

application.

Cross Reaction:

Peptide	%
β-Endorphin (human)	100
β-Endorphin (equine)	100
β-Endorphin (porcine)	100
β-Lipotropin	100
β-Endorphin (1-27) (human)	92
β-Endorphin (rat)	80
Ac-β-Endorphin (human)	70
Ac-β-Endorphin (1-27) (human)	50
β-Endorphin (6-31) (human)	43
β-Endorphin (camel, bovine, ovine)	39
β-Endorphin (1-26) (human)	37
Ac-β-Endorphin (1-26) (human)	30

Cross-Reactivity data is taken from Radioimmunoassay applications