


Anti-Crotonaldehyde (CRA) Monoclonal Antibody



Crotonaldehyde (CRA) is a representative carcinogenic aldehyde formed endogenously through lipid peroxidation. CRA is highly reactive aldehyde and reacts with lysine residue in protein. The reaction with CRA and lysine residue leads to the formation of numerous numbers of adducts. This antibody is specific for the CRA-modified protein.

- Catalog #:** N213630 (30 μ g of IgG)
- Source:** Mouse
- Immunogen:** CRA-modified keyhole-lympet hemocyanine.
- Subclass:** Mouse IgG_{2a}
- Application:** Immunohistochemistry. Recommended antibody concentration is 0.5-1.0 micro gram/mL on paraformaldehyde fixed tissue.
- Buffer Concentration:** 100 micro g/mL antibody in 10mM PBS containing 0.1 %NaN₃ and 0.5% BSA. Purified by Protein-A.
- Specificity:** Specific for CRA-modified protein (especially CRA-lysine adduct)
- Storage:** Less than -20°C
- Stability:** 6 months after date of receipt. For long term storage, aliquot product into individual tubes and freeze at -20 or -70°C. Avoid repeated freeze/thaw cycles.

For research use only, not for diagnostic use.

<Distributed by>

SCETI

DF Kasumigaseki Place, 3-6-7, Kasumigaseki,
Chiyoda-ku Tokyo 100-0013 Japan

URL: <http://www.sceti.co.jp/export/>

e-mail: exp-pet@sceti.co.jp

<Manufacturer>

Japan Institute For the Control of Aging (JaICA)
710-1, Haruoka, Fukuroi, Shizuoka 437-0122
JAPAN