

**anti-APP-C31 (C-terminal fragment of the caspase 3-cleaved APP) antibody, rabbit serum (ACT1)**

74-108 100 ul

The Alzheimer **Amyloid Precursor Protein** (APP) is a transmembrane protein whose abnormal processing is associated with the pathogenesis of Alzheimer's disease. APP695 lacking the protease inhibitor domain is the predominant form in neuronal tissues. APP695 is cleaved by caspases into the 664-residue amino (N)-terminal fragment that lacks the carboxyl C-terminal 31-residues (APPΔC31) and the 31-residues C-terminal fragment (**APP-C31**). Both fragments might be potent inducers of neuronal apoptosis. An antibody (named **ACT1**) against the N-terminus of caspase 3-generated APP C-terminal 31 aa of human APP695 (**APP-C31**) was raised in rabbit.

**Applications:**

1. Western blotting (dilution: 1/3,000-1/1,000)
2. Immunocytochemistry (dilution: 1/1,000-1/500)
3. ELISA

These applications were confirmed in the laboratory of Prof. K, Yoshikawa of Osaka University. (ref.3).

**Immunogen:** Synthetic peptide corresponding to the N-terminus of caspase 3-generated APP C-terminal 31 amino acids (aa 665-670 of human APP695).

**Specificity:** Reactive to human, mouse and rat. Specific to the N-terminal end of the caspase 3-generated APP-C31.

**Form:** Antiserum added with 0.05% sodium azide.

**Storage:** Shipped at 4 °C and stored at -20 °C.

**Data Link:** Swiss-Prot [P05067](#)

**References:** This antibody was used in ref.3.

1. Kang HG *et al.* (1987) "The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor." *Nature* **325**: 33-736 PMID: [2881207](#)
2. Selkoe DJ (1994) "Normal and abnormal biology of the beta-amyloid precursor protein." *Annu. Rev. Neurosci.* **17**: 489-517 PMID: [8210185](#)
3. Nishimura I *et al.* (2002) "Cell death induced by a caspase-cleaved transmembrane fragment of the Alzheimer amyloid precursor protein." *Cell Death Differ.* **9**: 199-208 PMID: [11840170](#)
4. Nishimura I *et al.* (2003) "Upregulation and antiapoptotic role of endogenous Alzheimer amyloid precursor protein in dorsal root ganglion neurons." *Exp. Cell Res.* **286**: 241-251 PMID: [12749853](#)

**Related products:** #74-102 anti-Activated Caspase3 antibody, #74-104 anti-APP (C-terminal) antibody,

< Distributed by > : **SCETI** SCETI K.K.



3-6-7 Kasumigaseki, Chiyoda-ku Tokyo 100-0013 JAPAN

Tel: +81-3-5510-2347 Fax: +81-3-5510-0134

E-mail: [exp-pet@sceti.co.jp](mailto:exp-pet@sceti.co.jp) URL: [www.sceti.co.jp/export/](http://www.sceti.co.jp/export/)

< Manufactured by > : BioAcademia, Inc. 7-7-18 Saito-Asagi, Ibaraki, Osaka 567-0085, JAPAN

To be continued.




Fig.1 Immunocytochemistry for APP $\Delta$ C31 and APP-C31: Generation of the caspase-cleaved fragments in NT2 neurons (neurally differentiated human NT2 embryonic carcinoma cells) overexpressing wild type APP (ref.3).

NT2 neurons were fixed 72 h after infection with adenovirus vector expressing wild-type APP and stained for the N-terminus of APP (P2-1, mouse monoclonal antibody), chromosomal DNA (Hoechst), the C-terminus of APP $\Delta$ C31 (SAC) or the N-terminus of APP-C31 (ACT1). Most of wild-type APP-accumulating neurons with shrunken and fragmented nuclei contain SAC- and ACT1-immunoreactivities (arrows), but non-neuronal cells are hardly labeled with SAC and ACT1 (arrowheads).

< Distributed by >: **SCETI** SCETI K.K.

**SCETI**

3-6-7 Kasumigaseki, Chiyoda-ku Tokyo 100-0013 JAPAN

Tel: +81-3-5510-2347 Fax: +81-3-5510-0134

E-mail: [exp-pet@sceti.co.jp](mailto:exp-pet@sceti.co.jp) URL: [www.sceti.co.jp/export/](http://www.sceti.co.jp/export/)

< Manufactured by >: BioAcademia, Inc. 7-7-18 Saito-Asagi, Ibaraki, Osaka 567-0085, JAPAN