

Anti SulfFP2/sulf-2 Polyclonal Antibody

Heparan sulfate and heparin play key roles in the binding of many growth and differentiation factors, and in signaling by other factors. In view of SulfFP's ability to modify the sulfation of heparan sulfate outside the cell, SulfFPs have been classified as new members of sulfatase family. The SulfFP gene is conserved in nematode, fruit fly and human.

SulfFP1 modifies the interaction between heparin binding proteins and the carbohydrate side-chain of heparan sulfate, and has a key role in regulating FGF and Wnt signaling. Changes in SulfFP1 levels in cancer cells have focused attention on SulfFP as a targeting molecule. SulfFP2 is a related enzyme resembling SulfFP1 in structure and activity.


Three SulfFP antibodies are available;

KM108: Specifically reacts with the rat SulfFP1

KM109: Specifically reacts with an N-terminal fragment of rat SulfFP2

KM110: Specifically reacts with a C-terminal fragment of rat SulfFP2

Package Size	25µg (100µL/vial)
Format	Rabbit polyclonal antibody (0.25mg/mL)
Buffer	PBS [containing 2% Block Ace as a stabilizer, 0.1%Proclin as a bacteriostat]
Storage	Store below -20°C Once thawed, store at 4°C. Repeated freeze-thaw cycles should be avoided.
Purification method	This antibody was established from the serum of a rabbit immunized with a N-terminal fragment of rat SulfFP2 (421-564 a.a.), expressed as a recombinant protein in E. coli. Purified by Protein G affinity chromatography.
Working dilution	For Western blotting: 1.0µg/ml For Immunocytochemistry: 8.0µg/ml



Western blotting

Sample: SulfFP2-transfected HEK293 cells supernatants

Immunocytochemistry

Sample: SulfFP2-transfected HEK293 cells

Preparation of antibodies and instruction:

Masu M. Nagamine S.

University of Tsukuba Graduate School of Comprehensive Human Sciences

Anti SulfFP2/sulf-2 Polyclonal Antibody**【Reference】**

1. Nagamine S. et al. :
Expression of a heparan sulfate remodeling enzyme, heparan sulfate 6-O-endosulfatase sulfatase FP2, in the rat nervous system.
Brain Res Dev Brain Res. 2005 Oct 6;159(2):135-43.
2. Morimoto-Tomita M. et al. :
Cloning and characterization of two extracellular heparin-degrading endosulfatases in mice and humans.
J Biol Chem. 2002 Dec 20;277(51):49175-85.
3. Ai X. et al. :
Substrate Specificity and Domain Functions of Extracellular Heparan Sulfate 6-O-Endosulfatases, QSulf1 and QSulf2.
J Biol Chem. 2006 Feb 24;281(8):4969-76.

Supplier**SCETI****SCETI K.K.**

3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013, JAPAN

Tel +81(3) 5510-2347 Fax +81(3) 5510-0133

URL: <http://www.sceti.jp/export/> e-mail: exp-pet@sceti.co.jp