

**Anti MARCKS Monoclonal Antibody
(Clone No. MAR11/2)**

Myristoylated alanine-rich C kinase substrate (MARCKS), a specific substrate for protein kinase C, is abundant in the brain. MARCKS binds calmodulin under Ca^{2+} or actin under non- Ca^{2+} conditions. Phosphorylation of MARCKS (P-MARCKS) can be used as an indicator of protein kinase C activation in intact cells. It has been shown that P-MARCKS has relatively weaker binding activity to calmodulin and actin compared with MARCKS.

Pathophysiologically, P-MARCS is also detected in the brain of Alzheimer's disease patients and in the smooth muscle cells of arteriosclerosis.


This antibody is specific to MARCKS and could be used for western blotting and immunohistochemistry.

Package Size 50µg (200µL/vial)
 Format Rat monoclonal antibody 0.25 mg/mL
 Buffer Block Ace as a stabilizer, containing 0.1% Proclin as a bacteriostat
 Storage Store below -20°C
 Once thawed, store at 4°C. Repeated freeze-thaw cycles should be avoided.

Clone No. MAR11/2
 Subclass IgG2a
 Purification method The lymphocytes from rat, immunized with C terminal peptides of rat MARCKS, were fused to mouse myeloma SP2 cells. The cell line (MAR11/2) with positive reaction was grown in serum free medium from which the antibody was purified by Protein G affinity chromatography.

Working dilution for immunohistochemistry: about 1~ 5µg/mL

The reactivity of anti MARCKS antibody using western blotting.



SCETI

**Anti MARCKS Monoclonal Antibody
(Clone No. MAR11/2)**

【Reference】

1. SD Rose et al, Differential expression of MARCKS and other calmodulin binding protein kinase C substrate in cultured neuroblastoma and glioma cells.
J.Neurochem.63, 2314-2323,1994

Supplier

SCETI
SCETI K.K.

3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013, JAPAN
Tel +81(3) 5510-2347 Fax +81(3) 5510-0133
URL: <http://www.sceti.jp/export/> e-mail: exp-pet@sceti.co.jp