

Anti human COUP-TF I mouse monoclonal antibody

COUP-TF I: Chicken ovalbumin upstream promoter-transcription factor I

Code No	PP-H8132-00 old No. ZZH8132H
Clone No.	H8132
Lot.	A-1
Concentration	1 mg/mL
Volume	100 uL
Ig Class	G2a

Description Chicken ovalbumin upstream promoter transcription factor I (COUP-TFI, EAR3, COUP-TFA; NR2F1) is a member of orphan nuclear receptor. COUP-TF I is expressed in specific regions of the rostral brain, in stripes in the presumptive hindbrain. COUP-TFI has varied roles in the development of the peripheral nervous system, such as early regionalization of the neocortex, differentiation of subplate neurons and guidance of thalamocortical axons. COUP-TFs were shown to interact with a number of other nuclear receptors.

Nomenclature NR2F1

Genbank X12795

Origin Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with Baculovirus-expressed recombinant human COUP-TF I (6-81 aa).

Specificity This antibody specifically recognizes human COUP-TF I and cross reacts with mouse and rat COUP-TF I. This antibody does not recognize human COUP-TF II and EAR2.

Purification Ammonium sulfate fractionation

Formulation Physiological saline with 0.1% Na₃N as a preservative.

Application / Recommended Concentration

In order to obtain the best results, optimal working dilutions should be determined by each individual user.

Western Blot 1 ug/mL

Non reducing Western Blot Not yet tested


ELISA 0.1 ug/mL

Immunoprecipitation Decide by use

Supershift Assay Not yet tested

Chromatin immunoprecipitation Not yet tested

Immunohistochemistry 10-50 ug/mL



Human Prostate gland paraffin section



Rat Cerebrum paraffin section

Storage Store at 2 - 8 °C up to one month. For long-term storage, the solution may be frozen in working aliquots. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended.

Reference Suh JM, *et al.* Mol Endocrinol, 2006, 20(12): 3412-20
Qin J, *et al.* Dev Dyn., 2007, 236(3): 810-20
Perilhou A, *et al.* Mol Cell Biol., 2008, 28(14): 4588-97

Notes Sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large amounts of water during disposal.

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

Not for Diagnostic or Therapeutic use. Purchase of this product does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written consent of Perseus Proteomics Inc. is prohibited.

MADE IN JAPAN

Jan 22, 2009

Distributed by

SCETI SCETI K.K.

3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013, Japan
Tel +81(3) 5510-2347 Fax +81(3) 5510-0133
URL: <http://www.sceti.co.jp/export/> e-mail: exp-pet@sceti.co.jp

Manufactured by

PPMX PERSEUS PROTEOMICS Perseus Proteomics Inc.

4-7-6 Komaba, Meguro-ku, Tokyo 153-0041, Japan
URL: <http://www.ppmx.com>