



## Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

---

### Monoclonal Antibody To Human Macrophages Possibly Recognizing A Scavenger Receptor

Monoclonal antibody X-4, together with X-14, and PM-2K, forms a particular group of macrophage specific antibodies which were tested at the Vth Leukocyte Typing Workshop held in Boston in 1993. The X-4 antigen was initially suspected to function as a scavenger receptor. However, experiments with CHO cells expressing Macrophage Scavenger Receptor (MSR) AI or AII showed that they were not recognized by this antibody.

---

<b>Product Number:</b>	T-1049
<b>Clone:</b>	X-4
<b>Host species, isotype:</b>	Mouse IgG1
<b>Quantity:</b>	100µg
<b>Format:</b>	Affinity purified, lyophilized  Reconstitute by adding 0.5ml distilled water. This stock solution contains 0.2mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), 10mg/ml bovine serum albumin (BSA) as a stabilizer and 0.01% thimerosal as a preservative.
<b>Stability:</b>	Original vial: 1 year at 4° - 8°C  Stock solution or aliquots thereof: 1 year at -20°C. Avoid repeated thawing and freezing.
<b>Applications:</b>	Tested for immunohistochemistry (IHC).  <b>Approximate working dilution for IHC:</b> Frozen sections: 2µg/ml (1:100) Paraffin sections: not tested.  Optimal dilutions should be determined by the end user. Suggested positive control: Human placenta.
<b>Immunogen:</b>	Cultured human monocytes.
<b>Antigen, epitope:</b>	The antigen is a protein of 100kDa as determined by immunoprecipitation. It is found on the cytoplasmic membranes of macrophages. The epitope has not been determined.



## Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

### Antigen distribution:

**Isolated cells:** Positive on peritoneal macrophages and on cultivated macrophages with serum supplementation. Positive in AML but not on blast cells. Negative under serum-free conditions, on freshly isolated monocytes and on fMLP activated neutrophils. U937 cells show increased antigen expression after TNF $\alpha$  and GM-CSF treatment.

**Tissue sections:** X-4 is positive on macrophages, Kupffer cells, mesangial cells and Hoffbauer cells. It is positive on intrafollicular and paracortical dendritic and spindle macrophages and with macrophages and foamy cells in *Xanthoma disseminatum* but negative in sarcoid granuloma and histiocytosis X.

### Specificity:

**Human:** Macrophages.

**Other species:** positive with rabbit, cat, dog, goat, pig, horse, monkey, weak on guinea pig, negative with bovine and rat.

---

### Comparison of Staining Patterns of X-4, X-14<sup>a</sup> and PM-2K<sup>b</sup> on different cell types

	X-4	X-14	PM-2K
Lymph nodes tingible body macrophages	$\pm$	-	$\pm$
Brain microglial cells	$\pm$	-	$\pm$
Blood monocytes 24h culture	$\pm$	-	-

+ = positive,  $\pm$  = weakly positive, - = negative

a: Product T-1050

b: Product T-1051

---

### Selected references

XU, H., STEINHAUSEN, F., PETERS, J.H.: Identification of Differentiation Antigens on Human Monocytes and Macrophages with Newly Produced Monoclonal Antibodies. *Exp. Cell. Biol.* 57, 122 (1989).

SHAW, S. et al. *Leukocyte Typing V: White Cell Differentiation Antigens*, Oxford University Press (1994) Ed. Schlossmann, S. et al. Abstracts M017, M030, M060, M081, M105, M132, M138, M145 (X-14 Workshop Codes = M51, BP264)

Zeng, L., M. Takeya, X. Ling, A. Nagasaki, K. Takahashi. : Interspecies reactivities of anti-human macrophage monoclonal antibodies to various animal species. *J Histochem Cytochem* 44: 845-853 (1996).

For in vitro research only. Caution: this product contains thimerosal, a poisonous and hazardous substance.