



Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

Biotinylated Monoclonal Antibody To Human IL-1Ra Interleukin-1 Receptor Antagonist

Monoclonal antibody 1384 is useful for studying biological effects of IL-1Ra *in vitro*. The interleukin-1 receptor antagonist (IL-1Ra) is a naturally occurring molecule that shares homology with IL-1 α and IL-1 β . It binds competitively to IL-1 receptors type I and type II with the same affinity as does IL-1, but does not induce any intracellular response. Two structural variants of IL-1Ra have been described: a 17kDa form that is secreted from monocytes, macrophages, neutrophils, and other cells (sIL-1Ra) and an 18-kDa form that remains in the cytoplasm of keratinocytes and other epithelial cells, monocytes, and fibroblasts (icIL-1Ra). An additional 16-kDa intracellular isoform of IL-1Ra has been described in neutrophils, monocytes, and hepatic cells. Both of the major isoforms of IL-1Ra are transcribed from the same gene through the use of alternative first exons. The production of IL-1Ra is stimulated by many substances including adherent IgG, other cytokines, and bacterial or viral components. The tissue distribution of IL-1Ra in mice indicates that sIL-1Ra is found predominantly in peripheral blood cells, lung, spleen, and liver, while icIL-1Ra is found in large amounts in skin. Studies in transgenic and knockout mice indicate that IL-1Ra is important in host defence against endotoxin-induced injury. IL-1Ra is produced by hepatic cells with the characteristics of an acute phase protein. Endogenous IL-1Ra is produced in numerous experimental animal models of disease as well as in human autoimmune and chronic inflammatory diseases. The use of neutralizing anti-IL-1Ra antibodies has demonstrated that endogenous IL-1Ra is an important natural antiinflammatory protein in arthritis, colitis, and granulomatous pulmonary disease.

Product Number:	T-1404
Clone:	1384.92
Host species, isotype:	Mouse IgG1
Quantity:	200 μ g
Format:	Affinity purified, biotinylated, lyophilized Reconstitute by adding 0.5ml distilled water. This stock solution contains 0.4mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), no stabilizer and 0.01% thimerosal as a preservative.
Stability:	Original vial: 1 year at 4° - 8°C Stock solution or aliquots thereof: 1 year at -20°C. Avoid repeated thawing and freezing.
Applications:	Tested for ELISA. Approximate working dilution: ELISA: 2 μ g/ml Optimal dilutions should be determined by the end user.
Immunogen:	Recombinant human IL-1Ra.
Antigen distribution:	The antigen is produced and secreted by monocytes and macrophages after stimulation with pro-inflammatory cytokines,



Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

LPS, and *in vitro* by immunoglobulins coated to surfaces after an induction phase of 4h. *In vitro* the quantity of IL-1Ra secreted depends on the dose and type of stimulus.

Specificity:

Human: IL-1 receptor antagonist; negative with human IL-1beta, IL-1 alpha.

Other: negative with mouse IL-1Ra, other species unknown.

Selected references

Towbin, H. et al., J Immunol Meth **170**:125-135 (1994)

Dinareello, C.A. and Thompson, R.C., Immunology Today **12**:404 (1991)

Dinareello, C.A. and Wolff, S.M., New Engl. J. Med. **328**:106 (1993).

For *in vitro* research only. Caution: this product contains thimerosal, a poisonous and hazardous substance.