

Peninsula Laboratories, LLC A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070 Tel: (800) 922-1516 • (650) 592-5392 Fax: (650) 595-4071 www.bachem.com

Monoclonal Antibody To Rat CD68 Rat Monocytes, Macrophages And Dendritic cells

Monoclonal Antibody ED1 is useful for detecting rat monocytes and macrophages and isolated dendritic (veiled) cells in the blood. The antibody recognises a single chain glycoprotein of 90-100kDa that is expressed predominantly on the lysosomal membrane of myeloid cells. Weak cell surface expression also occurs. The antigen is expressed by the majority of tissue macrophages and weakly by peripheral blood granulocytes. Studies have shown that the antigen recognised by ED1 has many characteristics in common with mouse macrosialin and human CD68.

Product Number:	T-3003		
Clone:	ED1		
Host species, isotype:	Mouse IgG1		
Quantity:	250µg		
Format:	Affinity purified, liquid		
	Supplied as 0.25ml solution. This stock solution contains 1mg/ml IgG, phosphate buffered saline pH 7.2 (PBS), 0.09% sodium azide as a preservative.		
Stability:	Stock solution or aliquots thereof: 1 year at -20°C. Avoid repeated thawing and freezing.		
Applications:	Tested for immunohistochemistry (IHC), has been described to work in Western Blotting and immunoprecipitation of the antigen, FACS (preferably on permeabilized cells).		
	Approximate working dilution for IHC: Frozen sections: 0.5-1µg/ml (1:1000 – 1:2000)		
	Paraffin sections:10µg/ml (1:100), no antigen retrieval required.		
	Paraffin sections:10µg/ml (1:100), no antigen retrieval required.		
Immunogen:	Paraffin sections:10µg/ml (1:100), no antigen retrieval required. Optimal dilutions should be determined by the end user.		
Immunogen: Antigen, epitope:	Paraffin sections:10µg/ml (1:100), no antigen retrieval required. Optimal dilutions should be determined by the end user. Suggested positive control: rat spleen.		
-	 Paraffin sections:10µg/ml (1:100), no antigen retrieval required. Optimal dilutions should be determined by the end user. Suggested positive control: rat spleen. Rat spleen cells CD68; ED1 recognises a 92kD cytoplasmic protein. The 		
Antigen, epitope:	 Paraffin sections:10µg/ml (1:100), no antigen retrieval required. Optimal dilutions should be determined by the end user. Suggested positive control: rat spleen. Rat spleen cells CD68; ED1 recognises a 92kD cytoplasmic protein. The epitope has not been further characterized. The antigen is found on 90% of monocytes in the peripheral blood. It is also expressed by 98% of isolated dendritic (veiled) 		



Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070 Tel: (800) 922-1516 • (650) 592-5392 Fax: (650) 595-4071 www.bachem.com

Distribution and staining pattern of macrophages identified by ED1, ED2 and ED3 in various organs (from Dijkstra et al., 1985, modified):

Monoclonal Antibody	ED1	ED2 EI	ED3	
Staining pattern	Granular, patchy cytoplasmic	Diffuse, membrane	Diffuse, membrane	
<u>Spleen</u>				
White pulp				
inner PALS	++	-	+ Weakly	
outer PALS	++	+	+ Weakly	
follicle	+/-	-	-	
marg. metallophils	+/- Weakly	-	+++ Branched	
marginal zone	+/- Weakly	-	+++ Branched	
Red Pulp	+++	+++	+++ Weakly	
<u>Lymph node</u> Cortex				
outer cortex	+/- Weakly	-	+++ Subsinusoidal	
branched	2			
paracortical area	++	+	-	
follicles	+/-	-	-	
Medulla	+++	+ 10-20%	+++	
Capsule	+	+	-	
Peyer`s patches				
Interfollicular area	+++	++	+ Small groups 3-4 cells	
Dome	+	-	-	
Follicle-	-	-		
Villi	+++ Apex	++ Apex basis	-	
Lung				
BALT	++	Periphery of BALT	-	
Perivascular/peribronchial	+	+++	-	
Alveolar	+++	-	-	
Thymus				
Cortex	++	++ Branched	-	
Medulla	++	-	-/+ Weakly	
Corticomedullary area	+++	+++	-	
Capsule	+++ Branched	+++ Branched	++ Branched	
<u>Liver</u>	+++ Branched	+++ Branched	++ Branched	
Bone marrow	+++ Monocytes/ macrophages	++ Macrophages	-	

++ = A considerable number stained + = Few stained -/+ = Very few stained or none at all

Selected references

DIJKSTRA, C.D et al.: The heterogeneity of mononuclear phagocytes in lymphoid organs: distinct macrophage subpopulations in the rat recognised by monoclonal antibodies ED1, ED2 and ED3. Immunology: **54**, 589 - 599 (1985).

BEELEN, R.H.J et al.: Monoclonal Antibodies ED1, ED2, and ED3 Against Rat Macrophages: Expression of Recognized Antigens in Different Stages of Differentiation. Transplantation Proceedings: **XIX**, (3), 3166-3170 (1987).

DAMOISEAUX, J.G.M.C. et al.: Rat macrophage lysosomal mambrane antigen recognised by monoclonal antibody ED1. Immunology **83**, 140-147 (1994)



Peninsula Laboratories, LLC A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070 Tel: (800) 922-1516 • (650) 592-5392 Fax: (650) 595-4071 www.bachem.com

BAUER, J. et al.: Phagocytic activity of macrophages and microglial cells during the course of acute and chronic relapsing experimental autoimmune encephalomyelitis. J. Neurosci. Res. **38**, 365-375 (1994).

For in vitro research only. Caution: this product contains sodium azide, a poisonous and hazardous substance.