

Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070 Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071 www.bachem.com

Biotinylated Monoclonal Antibody to Rat Granulocytes

Monoclonal antibody RK-4 detects all rat granulocytes in frozen, and in paraffin embedded rat tissue sections after enzyme digestion.

Product Number: T-3119

Clone: RK-4

Host species, isotype: Mouse IgG1

Quantity: 150μg

Format: Affinity purified, biotinylated, lyophilized

Reconstitute by adding 0.5ml distilled water. This stock solution contains 0.3mg/ml lgG, phosphate buffered saline pH 7.2 (PBS), 10mg/ml bovine serum albumin (BSA) as a stabilizer

and 0.01% thimerosal as a preservative.

Stability: Original vial: 1 year at 4° - 8°C

Stock solution or aliquots thereof: 1 year at -20°C. Avoid

repeated thawing and freezing.

Applications: Tested for immunohistochemistry (IHC); has been described to

work in FACS.

Approximate working dilution for IHC:

Frozen sections: 0.75µg/ml (1:400)

Paraffin sections: does not react on routinely processed

paraffin sections

Optimal dilutions should be determined by the end user. Suggested positive control: Rat spleen, blood smears.

Immunogen: Peritoneal cells.

Antigen, **epitope**: The antigen is localized on the cell surface. The epitope has

not been further characterized.

Antigen distribution: Isolated cells Only a low percentage of cells from peripheral

blood stains positive. Positive on peritoneal cells 18h after thioglycollate injection (95% granulocytes). Negative on peritoneal cells 4 days after thioglycollate injection.

Negative on isolated lymphocytes, monocytes, erythrocytes

and platelets.



Peninsula Laboratories, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070 Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071 www.bachem.com

Specificity: Rat: pan-granulocytes,.

Other species: human: negative, other species not tested.

Tissue staining:

RK-4 reaction pattern:

Tissues tested	Results
Bone marrow	positive on granulocytes
	positive on granulocyte precursors positive on segmented cells
	positive on band cells
	positive on metamyelocytes
	positive on myelocytes
Lung	negative on alveolar macrophages; interstitial tissues are occasionally positive.
Kidney	negative; on perivascular granulocytes occasionally positive.
Skin	negative; on perivascular granulocytes occasionally positive.
Liver	negative; on perivascular granulocytes occasionally positive.
Blood vessel wall	negative

Selected references

STEINHAUSEN: F;, H. Xu, H.-J. Groene, H.J. Peters: A monoclonal Antibody Selectively Recognizing Rat Granulocytes. Immunobiol: 181: 324 (1990)

For in vitro research only. Caution: this product contains thimerosal, a poisonous and hazardous substance.