

# PENINSULA LABORATORIES, LLC

## A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070 Tel: (800) 922-1516 • (650) 592-5392 Fax: (650) 595-4071

www.bachem.com

### Cat. No. T-4146

## Rabbit anti-Amylin (mouse, rat)

## **Specifications:**

Preparation: This polyclonal antibody was collected from rabbits immunized

with a synthetic peptide as the immunogen.

Amount: Each vial contains 50 ul of whole serum in lyophilized form.

Reconstitute with 50 ul of distilled water for the equivalent of

undiluted antiserum.

Storage: This antibody is supplied as a lyophilized powder. Store at 2° to

8°C upon receipt. For long-term storage, store lyophilized powder at -20°C. Once reconstituted, the materials should be stored frozen in aliquot at -20°C to -70°C. **Avoid repeated** 

freeze/thaw cycles.

Applications: Immunohistochemistry, Immunocytochemistry, and other

applications.

Sample Types: The antibody can react with a variety of sample preparation

including

frozen tissue sections, paraffin embedded tissue sections and fixed

cell cultures.

### Recommended starting dilution for Immunohistochemical staining:

Immunofluorescence: 1:250 to 1:16000

PAP or ABC: 1:250 to 1:16000 LAB-SA: 1:250 to 1:16000

It is advised for each laboratory to determine optimal dilutions for a specific

application.

#### Cross Reaction:

| Peptide                                  | %    |
|--|------|
| Amylin (mouse, rat)                      | 100  |
| Amylin (cat)                             | 100  |
| Amylin (human)                           | 100  |
| α-CGRP (human)                           | 0.01 |
| α-CGRP (rat)                             | 0.01 |
| β-CGRP (human)                           | 0.01 |
| β-CGRP (rat)                             | 0.01 |
| Glucagon (1-29) (human, bovine, porcine) | 0    |
| Insulin (human)                          | 0    |
| NPY (porcine)                            | 0    |
| VIP (human, bovine, porcine, rat)        | 0    |
| Pancreatic Polypeptide (human)           | 0    |

Cross-Reactivity data is taken from Radioimmunoassay applications