



PENINSULA LABORATORIES, LLC

A Member of the Bachem Group

305 Old County Road, San Carlos, CA 94070

Tel: (800) 922-1516 • (650) 592-5392

Fax: (650) 595-4071

www.bachem.com

Cat. No. T-4170

Rabbit anti- β -Atrial Natriuretic Factor (1-28) (Dimer, Antiparallel) (human)

Specifications:

- Preparation:** This polyclonal antibody was collected from rabbits immunized with a synthetic peptide as the immunogen.
- Amount:** Each vial contains 50 μ l of whole serum in lyophilized form. Reconstitute with 50 μ l of distilled water for the equivalent of undiluted antiserum.
- Storage:** This antibody is supplied as a lyophilized powder. Store at 2° to 8°C upon receipt. For long-term storage, store lyophilized powder at -20°C. Once reconstituted, the materials should be stored frozen in aliquot at -20°C to -70°C. **Avoid repeated freeze/thaw cycles.**
- Applications:** Immunohistochemistry, Immunocytochemistry, and other applications.
- Sample Types:** The antibody can react with a variety of sample preparation including frozen tissue sections, paraffin embedded tissue sections and fixed cell cultures.

Recommended starting dilution for Immunohistochemical staining:

Immunofluorescence: 1:250 to 1:16000

PAP or ABC: 1:250 to 1:16000

LAB-SA: 1:250 to 1:16000

It is advised for each laboratory to determine optimal dilutions for a specific application.

Cross Reaction:

Peptide	%
β -Atrial Natriuretic Factor (1-28) (Dimer, Antiparallel) (human)	100
ANF (4-28) (human)	3.4
ANF (1-28) (human)	2.9
Urodilantin	2.8
ANF (rat)	0
ANP (1-11) (human)	0
BNP-32 (porcine)	0
CNP (32-53) (human, porcine, rat)	0

Cross-Reactivity data is taken from Radioimmunoassay applications