

Anti basic FGF (1-9) (Human) Serum

Cat. No. Y251

Lot No. 06280703

Description: This antiserum was raised in a rabbit by immunization with a bovine serum albumin (BSA) conjugate of synthetic basic FGF (1-9)-Tyr (human) peptide. The product vial contains 50 μ L of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with N-terminal portion of basic FGF(human).

Immunogen: Synthetic basic FGF (1-9)-Tyr-(human)-BSA conjugate

Host: Rabbit

Amino Acid Sequence of basic FGF (1-9)(human)¹⁾:

1 9
PALPEDGGS

Product Form: Lyophilized unpurified serum

Size: 50 μ L

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).

Storage: The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

Suggested Working Dilution Range: 1:750 (final dilution ~1:6,000) for radioimmunoassay; 1: 200- 1,000 for immunohistochemistry (frozen section). Optimal dilution should be determined by each laboratory for each application.

Specificity (based on radioimmunoassay): Recombinant basic FGF (human) 100%, basic FGF (1-9)-Tyr (human) 100%

Positive Control (immunohistochemistry): Human pituitary gland

REFERENCES:

1) J.R. Abraham, J.L. Whang et al., Human basic fibroblast growth factor: nucleotide sequence and genomic organization. EMBO Journal 5:2523-2529, 1986

FOR RESEARCH USE ONLY

<Distributed by>

SCETI K.K.

DF Kasumigaseki Place, 3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan

URL: <http://www.sceti.co.jp/export/> e-mail: exp-pet@sceti.co.jp

<Manufacturer>

Yanaihara Institute Inc.

2480-1 Awakura, Fujinomiya-shi, Shizuoka 418-0011 JAPAN