

Anti Chromogranin A (344-374) (Human) Serum

Cat. No. Y293 Lot No. 7670424

Description: This antiserum was raised in a rabbit by immunization with a carrier free synthetic chromogranin A (344-374) (human) peptide. The product vial contains 50 μ L of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with chromogranin A related peptides.

Immunogen: Synthetic chromogranin A (344-374) (human), carrier free

Host: Rabbit

Amino Acid Sequence of Chromogranin A (344-374) (human)¹⁾:

344

374

EEEEDNR DSSMKLSFRA RAYGFRGPGP OLRR

Product Form: Lyophilized unpurified serum

Size: 50 μ L

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN_3 0.1%).

Storage: The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

Suggested Working Dilution Range: 1:500-1,200 (final dilution ~1:3,500-8,400) for radioimmunoassay^{2,3)}; 1:2,500 for enzyme immunoassay⁴⁾, 1: 1,000 for immunohistochemistry (frozen section). Optimal dilution should be determined by each laboratory for each application.

Specificity (based on radioimmunoassay): CgA (344-374) (human) 100%, CgA (359-389) (rat) 33.5%, not react with CgA (375-389) (rat) and CgA (360-374) (human)

Positive Control (immunohistochemistry): Rat pituitary gland and adrenal gland

Species Tested: Human

REFERENCES:

- 1) D.S. Konecki, U.M. Benedum et al., The primary structure of human chromogranin A and pancreastatin. *Journal of Biological Chemistry*, 262:17026, 1987
- 2) Y. Nishikawa, J. Li et al., Region-specific radioimmunoassay for human chromogranin A, *Biomedical Research*, 19:245-252, 1998
- 3) H. Yanaihara, M. Hata et al., Application of region-specific immunoassay for human chromogranin A: substantial clue for detection and measurement of chromogranin A in human plasma, *regulated Peptide*, 80:83-90, 1999
- 4) S. Nagasawa, Y. Nishikawa et al., Simple enzyme immunoassay for the measurement of immunoreactive chromogranin A in human plasma, urine, and saliva, *Biomedical Research*, 19: 407-410, 1998

FOR RESEARCH USE ONLY

<Distributed by>

SCETI K.K.

DF Kasumigaseki Place, 3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013 Japan

URL: <http://www.sceti.co.jp/export/> e-mail: exp-pet@sceti.co.jp

<Manufacturer>

Yanaihara Institute Inc.

2480-1 Awakura, Fujinomiya-shi, Shizuoka 418-0011 JAPAN