

T4 DNA Ligase

02-050 20,000 U (400U/μl), 02-050-5 5 X 20,000 U (400U/μl)

Bacteriophage T4 derived DNA ligase catalyzes the formation of phosphodiester bonds between 3'-OH termini and 5'-P termini in duplex DNA or RNA (1). This enzyme will join blunt end and cohesive end termini as well as repair single stranded nicks in duplex DNA, RNA or DNA/RNA hybrids.

T4 DNA ligase was expressed in *E.coli* in large quantities and highly purified. MW is 55.3 kDa.

Applications:

- 1) Insertion of DNA fragment into a vector
- 2) Linker (or Adaptor) ligation with DNA fragment

Storage conditions:

10mM Tris-HCl (pH 7.6), 50mM KCl, 0.1mM EDTA, 1mM dithiothreitol, 50% glycerol

Store at -20

Concentration:

400 U/μl, where one unit is the amount of enzyme that ligates more than 90% of 6 μg of λDNA-HindIII fragments in a 20μl mixture in 30 minutes at 16 °C.

Quality Assurance:

Greater than 95% protein determined by SDS-PAGE (CBB staining)

The absence of endonucleases and exonucleases was confirmed.



Reagents Supplied with Enzyme:

10 x T4 Ligase Reaction Buffer (T4-Lig): 500mM Tris-HCl (pH 7.6), 100mM MgCl₂, 10 mM ATP, 100mM dithiothreitol

Data Link: Swiss-Prot [P00970](#)

References:

1. Weiss, B. *et al.* (1968) "Enzymatic breakage and joining of deoxyribonucleic acid." *J. Biol. Chem.* **243**: 4543-4555 PMID: [4879167](#)



SCETI

SCETI K.K.

3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013, JAPAN
Tel +81(3) 5510-2347 Fax +81(3) 5510-0133
URL: <http://www.sceti.jp/export/> e-mail: exp-pet@sceti.co.jp



Fig.1 SDS-PAGE of T4 DNA ligase protein

Fig.2 DNA ligation activity

Ligation of Hind III fragments of λ DNA using 1 unit of T4 DNA ligase

Incubation at 16 °C for 0, 10, 20, 30, and 60 min.