

## **Anti-Dnmt1 (1-248) antibody, affinity-purified (rabbit polyclonal)**

70-201

50  $\mu$ g

Dnmt1 (DNA (cytosine-5)-methyltransferase 1) has a role in the establishment and regulation of tissue-specific patterns of methylated cytosine residues (epigenetics). Hypermethylation in promoter regions are associated with repression of the genes. Aberrant methylation patterns are associated with certain human tumors and developmental abnormalities. This antibody was raised by Prof. S. Tajima of Osaka Univ. and used in Ref.3

### **Applications**

- 1) Western blotting (0.2~1  $\mu$ g /ml)
- 2) Immunoprecipitation. Especially good for native form of Dnmt1 and useful for analysis of the protein complex containing Dnmt1
- 3) Indirect immuno-fluorescence staining: Excellent

Other applications were not tested

### **Properties of the product**

**Antigen:** Highly purified recombinant mouse Dnmt1 (amino acids 1-248), soluble form

**Antibody:** Affinity-purified with the recombinant Dnmt1

**Reactivity:** Mouse and human. Not tested with other species

**Form:** 1 mg/ml in PBS, 50% glycerol, 0.05% sodium azide (and trace of ammonium sulfate)

**Storage:** -20°C (long period; -80°C)

**Reference:** This antibody was used for western blotting and immuno-fluorescence staining in Ref. 3.

1. Di Croce L, *et al.* (2002). Methyltransferase recruitment and DNA hypermethylation of target promoters by an oncogenic transcription factor. *Science* 295: 1079-82.
2. Rhee I, *et al.* (2002). DNMT1 and DNMT3b cooperate to silence genes in human cancer cells. *Nature* 416: 552-6.
3. Sharif K, Muto M, Takebayashi S, *et al.* (2007) The SRA protein Np95 mediates epigenetic inheritance by recruiting Dnmt1 to methylated DNA. *Nature* 450:908-12.

**The data of western blotting and immuno-fluorescence staining obtained by using this antibody are shown in the next page**


**Related product: # 70-203 Anti-Dnmt1 (1037-1086) antibody, affinity-purified (rabbit polyclonal)**

**Especially good for immunoprecipitation of denatured Dnmt1 and useful for ChIP assays**

**Cross-reacts with human, mouse and xenopus Dnmt1**

<Distributed by >: SCETI K.K. DF Kasumigaseki Place, 3-6-7 Kasumigaseki,  
Chiyoda-ku Tokyo 100-0013 JAPAN Tel: +81-3-5510-2347 Fax: +81-3-5510-0134  
E-mail: [exp-pet@sceti.co.jp](mailto:exp-pet@sceti.co.jp) URL: [www.sceti.co.jp/export/](http://www.sceti.co.jp/export/)

<Manufactured by> : BioAcademia, Inc. 7-7-18 Saito-Asagi, Ibaraki, Osaka 567-0085, JAPAN



**Figure. Use of anti-Dnmt1 (1-248) antibody for western blotting and immuno-fluorescence staining**

**a**, Subnuclear localization of Dnmt1 (green), Np95 (red) and PcnA (blue) in mouse embryonic stem cells (E14) during cell-cycle progression (bottom). Merged images for Np95 and Dnmt1 are shown. Profile of DNA content in exponentially growing ECSs is shown (top)

**b**, Association of Np95 with Dnmt1 and PcnA in HeLa cell nuclear extracts. Human NP95 was tagged by biotin-binding domain and stably expressed in HeLa cells together with the *E. coli BirA* biotin ligase gene. Biotylated Np95 was captured by streptavidin beads. The captured proteins were eluted from the beads and analyzed by western blotting using antibodies against Dnmt1 and PcnA.

**c**, Catalytic activity of NP95 associated Dnmt1 in HeLa cells. The presence of Dnmt1 in the NP95 complexes was confirmed by western blot and the DNA methyltransferase activity of the extracts was measured

(The data were kindly provided by Dr. M. Muto of Riken Research Center for Allergy and Immunology)