

KO454 For research use only

Anti Mouse Rim1 Polyclonal Antibody

This antibody was prepared by Dr. Yasuo Mori, Kyoto University.

Code No. KO454
Terget Rim1

Category TRP channel
Gene ID 116837
Primary Source MGI:2152971

Synonyms Rim; RIM1; RIM1a; Serg1; Rab3ip1; KIAA0340; RIM1alpha;

mKIAA0340; C030033M19Rik; Rims1

Type Polyclonal Antibody

Immunogen Partial peptide of Mouse Rim1 N-terminal region

Raised in Rabbit Myeloma -

Clone number -

Purification Antigen Affinity
Source Rabbit Serum

Isotype Cross Reactivity -

LabelUnlabeledConcentration0.25 mg/mL

Contents (Volume) 25 µg (100 µL/vial)

Buffer PBS [containing 2% Block Ace as a stabilizer, 0.1% Proclin

as a bacteriostat]

Storage Store below -20°C. Once thawed, store at 4°C. Repeated

freeze-thaw cycles should be avoided.

Application ELISA,WB,IP

| ELISA | WB | IHC | ICC |
|-------|------------|------------|----------------|
| 1.0 | 1.0 | Not tested | Not tested |
| IP | FCM | IF | Neutralization |
| 5.0 | Not tested | Not tested | Not tested |

(µg/mL)

Reference

- 1. Kiyonaka S, et al. RIM1 confers sustained activity and neurotransmitter vesicle anchoring to presynaptic Ca2+ channels. Nat Neurosci. 2007 Jun;10(6):691-701. *Application Reference
- 2. Schoch S, et al. RIM1alpha forms a protein scaffold for regulating neurotransmitter release at the active zone. Nature. 2002 Jan 17;415(6869):321-6.

UniPlot Summary

//Function: Rab effector involved in exocytosis. May act as scaffold protein that regulates neurotransmitter release at the active zone. Essential for maintaining normal probability of neurotransmitter release and for regulating release during short-term synaptic plasticity. //Sequence similarities: Contains 2 C2 domains. Contains 1 FYVE-type zinc finger. Contains 1 PDZ (DHR) domain. Contains 1 RabBD (Rab-binding) domain.

