

freeze-thaw cycles.

For research use only

KO617 Anti mouse AIM Monoclonal Antibody

Clone No. 23B12

Target	mouse AIM
Category	immunology
Gene ID	11801
Primary Source	MGI:1334419
Synonyms	CD5L, AAC-11, AIM/Spalpha, Api6, Pdp 1/6, Sp-alpha
Туре	Monoclonal Antibody
Immunogen	recombinant mouse AIM
Raised in	Wistar Rat
Myeloma	P3U1
Clone number	23B12 (#36)
Purification	ProteinG
Source	Serum-free medium
lsotype	lgG2a
Cross Reactivity	Not tested
Label	Unlabeled
Concentration	-
Contents (Volume)	-
Buffer	PBS
Storage	Store at - 20°C long term, store at 4°C short term. Avoid repeated t

Application

ELISA, WB, IHC, ICC, IP

ELISA	WB	IHC	ICC
1.0	1.0	10.0	10.0
IP	FCM	IF	Neutralization
5.0	Not tested	Not tested	-
			(µg/mL)

Reference

Miyazaki T et al. AlMing at Metabolic Syndrome- Towards the Development of Novel Therapies for Metabolic Diseases via Apoptosis Inhibitor of Macrophage (AIM) -Circ. J., 2011, 75, 2522-2531

Kurokawa et al. Apoptosis inhibitor of macrophage (AIM) is required for obesity-associated recruitment of inflammatory macrophages into adipose tissue. Proc Natl Acad Sci USA 2011, 108, 12072-12077

Kurokawa et al. Macrophage-derived AIM is endocytosed into adipocytes and decreases lipid droplets via inhibition of fatty acid synthase activity. Cell Metab. 2010, 11, 479-492

UniProt Summary

//Function: May play a role in the regulation of the immune system. Seems to play a role as an inhibitor of apoptosis.

//Subcellular location:Secreted.

//Tissue specificity: Expressed in thymus, liver, spleen and lymph nodes.

//Post-translational modification: Glycosylated.

//Sequence similarities: Contains 3 SRCR domains.

C Distributed by SCETI K.K.
SCETI K.K.
3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013, Japan

Tel: +81(3) 5510-2652 Fax: +81(3) 5510-0133

URL : http://www.sceti.co.jp/export E-mail : medical@sceti.co.jp