

## Anti human TR4 mouse monoclonal antibody


TR4: Tasticular receptor 4

Code No	PP-H0107B-00
Clone No.	H0107B
Lot.	A-1
Concentration	1 mg/mL
Volume	100 uL
Ig Class	G2a
Description	TR4(TAK1, NR2C2) is a member of orphan nuclear receptor. TR4 was originally cloned from lymphoblastoma Raji cells or mouse brain cDNA library. No ligand has been reported. Northern blot shows TR4 is transcribed as a 9kb mRNA in many tissues and as a 2.8kb mRNA in testis, mainly in spermatocytes. TR4 has two isoforms called TR4alpha1 and TR4 alpha2, which differ in 19 amino acids coded by two separate exons. Both products translated from 9kb transcript are ubiquitously expressed. Since TR4 binds to the same elements for the RAR-RXR or TR-RXR heterodimers, TR4 may have an inhibitory effect for retinoic-acid mediated transactivation.
Nomenclature	NR2C2
Genbank	L27586
Origin	Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with Baculovirus-expressed recombinant human TR4 (23-52 aa).
Specificity	This antibody specifically recognizes human TR4 and cross reacts with mouse and rat TR4.
Purification	Ammonium sulfate fractionation
Formulation	Physiological saline with 0.1% NaN <sub>3</sub> as a preservative.

### Application / Recommended Concentration

In order to obtain the best results, optimal working dilutions should be determined by each individual user.

Western Blot	2 ug/mL
Non reducing Western Blot	Not yet tested
ELISA	0.1 ug/mL (A450=0.2)
Immunoprecipitation	Decide by use
Supershift Assay	Not yet tested
Chromatin immunoprecipitation	Not yet tested
Immunohistochemistry	10µg/mL



Rat cerebellum  
Purkinje cells and granular cells

Rat testis  
primary and secondary  
spermatocytes

Storage	Store at 2 - 8 °C up to one month. For long-term storage, the solution may be frozen in working aliquots. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended.
---------	---

### Reference

Notes	Sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large amounts of water during disposal.
-------	---

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

Not for Diagnostic or Therapeutic use. Purchase of this product does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written consent of Perseus Proteomics Inc. is prohibited.

**MADE IN JAPAN**

Aug 29, 2006

Distributed by

**SCETI** SCETI K.K.

3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013, Japan  
Tel +81(3) 5510-2347 Fax +81(3) 5510-0133  
URL: <http://www.sceti.co.jp/export/> e-mail: [exp-pet@sceti.co.jp](mailto:exp-pet@sceti.co.jp)

Manufactured by

**PPMX**  
PERSEUS PROTEOMICS

Perseus Proteomics Inc.

4-7-6 Komaba, Meguro-ku, Tokyo 153-0041, Japan  
URL: <http://www.ppmx.com>