

Anti human PXR common mouse monoclonal antibody

PXR: Pregnane X receptor, SXR

	PP-H4417-00	Application / Recommended Concentration In order to obtain the best results, optimal working dilutions should be determined by each individual user.		
Code No				
Clone No.	H4417	Westerr	n Blot	1ug/mL
Lot.	A-1	Non red	ucing Western Blot	3ug/mL
Concentration	1 mg/mL			
Volume	100 uL	ELISA		2ug/mL (A450=1.0)
Ig Class	G2a	Immunc	precipitation	Decide by use
Description	Pregnane-activated receptor (PXR,SXR, PAR, PAR1, PAR2, NR112) is a member of nuclear receptor subfamily , which each orthologues historically given	Superst	ift Assay	Not yet tested
	different names as pregnane-activated receptor in mice and steroid- and xenobiotic-sensing nuclear receptor in human, PXR binds to rifampicin (an antibiotics) is the most efficient activator in human. Various studies	Chromatin immunoprecipitation Not yet tested		
	revealed PXR regulates CYP3A gene expression as well as other xenobiotic metabolisms, such as oxidation, conjugation and transport. Many chemicals are known to bind for PXR as activators, eg. the HIV protease inhibitor ritonavir, the anticancer drug paclitaxel, the endocrine disruptor bisphenol A. Expression of PXR founds in the liver, small intestine and colon in the human, rabbit and mouse where CYP3A genes are expressed or induced.	Immunohistochemistry Not yet tested		
Nomenclature	NR1I 2			
Genbank	AF084645			
Origin	Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with Baculovirus-expressed recombinant human PXR-1(1-40 aa).	Storage	the solution may be Repeated freezing a	to one month. For long-term storage, frozen in working aliquots. and thawing is not recommended. be freezer is not recommended.
Specificity	This antibody specifically recognizes human PXR-1 and PXR-2. Not yet tested in other species.	Reference		
Purification	Ammonium sulfate fractionation	Notes	Sodium azide may r	eact with lead and copper plumbing
Formulation	Physiological saline with 0.1% NaN3 as a preservative.		to form explosive metal azides. Flush with large amounts of water during disposal.	

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

SCETI K.K.

3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013, Japan Tel +81(3) 5510-2347 Fax +81(3) 5510-0133

URL: http://www.sceti.co.jp/export/ e-mail: exp-pet@sceti.co.jp

Distributed by

SCE'

Not for Diagnostic or Therapeutic use. Purchase of this product does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written consent of Perseus Proteomics Inc. is prohibited.

MADE IN JAPAN

Aug 24, 2006



4-7-6 Komaba, Meguro-ku, Tokyo 153-0041, Japan URL: http://www.ppmx.com