

Anti human COUP-TF II mouse monoclonal antibody

COUP-TF II: Chicken ovalbumin upstream promoter-transcription factor II

Code No	PP-H7147-10
Clone No.	H7147
Batch	A1F1
Size	100ug lyophilized antibody
Formulation	Lyophilized from phosphate-buffered saline (PBS) containing 5% Trehalose as a stabilizer.
Ig Class	G2a
Nomenclature	NR2F2
Genbank	M64497
Description	Chicken ovalbumin upstream promoter transcription factor II (COUP-TFII, ARP-1, COUP-TFB; NR2F2) is a member of the nuclear orphan receptor family. COUP-TFII is expressed in tongue, follicles of vibrissae, cochlea and in stroma of nasal septum. COUP-TFII has roles in angiogenesis, vascular remodeling and heart development. COUP-TFs were shown to interact with a number of other nuclear receptors.
Origin	Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with Baculovirus-expressed recombinant human COUP-TF II (43-64 aa).
Specificity	This antibody specifically recognizes human COUP-TF II and cross reacts with mouse and rat COUP-TF II. This antibody does not recognize human COUP-TF I and EAR2.
Purification	Ammonium sulfate fractionation.

Application / Recommended Concentration

In order to obtain the best results, optimal working dilutions should be determined by each individual user.

Western Blot 1 ug/mL

Non-reducing Western Blot Not yet tested


ELISA 0.1 ug/mL

Immunoprecipitation Decide by use

Supershift Assay Not yet tested

Chromatin immunoprecipitation Not yet tested

Immunohistochemistry 10 ug/mL



Human
Convoluted tubule
paraffin section

Rat
Glomerular
paraffin section

Reconstitution Reconstitute with desired volume of distilled water. Reconstitution in 0.1 ml of distilled water yields an antibody concentration of 1mg/ml. Please note reconstituted solution will contain original PBS and 5% Treharose (see Formulation).

Storage Lyophilized samples are stable for twelve months when held at 2 - 8 C. Upon reconstitution, the antibody can be stored at 2 - 8 C up to one month. For long-term storage, the solution may be frozen in working aliquots. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended.

Reference Lee CT, *et al.* Mol Cell Biol., 2004, 24(24): 10835-43
You LR, *et al.* Proc Natl Acad Sci USA. 2005, 102 (45):16351-6
Suh JM, *et al.* Mol Endocrinol, 2006, 20(12): 3412-20
Qin J, *et al.* Dev Dyn., 2007, 236(3): 810-20
Perilhou A, *et al.* Mol Cell Biol., 2008, 28(14): 4588-97
Li L, *et al.* Cell Metab., 2009, 9(1): 77-87

Notes

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

Not for Diagnostic or Therapeutic use. Purchase of this product does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written consent of Perseus Proteomics Inc. is prohibited.

MADE IN JAPAN

Jan 22, 2009

Distributed by

SCETI SCETI K.K.

3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013, Japan
Tel +81(3) 5510-2347 Fax +81(3) 5510-0133
URL: <http://www.sceti.co.jp/export/> e-mail: exp-pet@sceti.co.jp

Manufactured by

PPMX
PERSEUS PROTEOMICS

Perseus Proteomics Inc.

4-7-6 Komaba, Meguro-ku, Tokyo 153-0041, Japan
URL: <http://www.ppmx.com>