

Anti human PPAR gamma common mouse monoclonal antibody

PPAR gamma: Peroxisome Proliferator-Activated Receptor gamma

Code No	PP-K8713-00 old No. 2ZK8713H
Clone No.	K8713
Lot.	A-2
Concentration	1 mg/mL
Volume	100 uL
Ig Class	G2a
Description	Peroxisome proliferator-activated receptor gamma (PPARγ; NR1C3) is a member of orphan nuclear receptor. Oxidized metabolites of linoleic acid, 9-hydroxyoctadecenoic acid (9-HODE) and 13-HODE are activators and ligands of PPARγ. PPARγ is expressed in white adipose tissue, intestinal mucosa, colon, spleen, monocytes, macrophages, retina, cartilage, osteoclast and skeletal muscle. PPARγ plays important roles in lipid and glucose metabolism, and have been implicated in obesity-related metabolic diseases such as hyperlipidemia, insulin resistance, and coronary artery disease. Three members were called PPARα, β, γ. Three N-terminal isoforms, called γ1, γ2 and γ3, are known to arise by alternative splicing and promoter usage from the PPARγ gene. RXR is an obligate partner for PPAR.
Nomenclature	NR1C3
Genbank	U79012
Origin	Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with Baculovirus-expressed recombinant human PPAR gamma2 (2-136 aa) .
Specificity	This antibody specifically recognizes human PPAR gamma1 and 2, and cross reacts with mouse PPAR gamma1 and 2. This antibody does not recognize human PPAR alpha and delta. Not yet tested in other species.
Purification	Ammonium sulfate fractionation
Formulation	Physiological saline with 0.1% NaN ₃ as a preservative.

Application / Recommended Concentration

In order to obtain the best results, optimal working dilutions should be determined by each individual user.

Western Blot	2 ug/mL
Non reducing Western Blot	Not yet tested
ELISA	0.1 ug/mL
Immunoprecipitation	Decide by use
Supershift Assay	100 ug/mL
Chromatin immunoprecipitation	Decide by use
Immunohistochemistry	Not yet tested

Storage Store at 2 - 8 °C up to one month. For long-term storage, the solution may be frozen in working aliquots. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended.

Reference Tanaka T, *et al.* J Atheroscler Thromb, 9(5): 233 -241, 2002.

Notes Sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large amounts of water during disposal.

FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

Not for Diagnostic or Therapeutic use. Purchase of this product does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written consent of Perseus Proteomics Inc. is prohibited.

MADE IN JAPAN

Aug 23, 2006

Distributed by

SCETI SCETI K.K.

3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013, Japan
Tel +81(3) 5510-2347 Fax +81(3) 5510-0133
URL: <http://www.sceti.co.jp/export/> e-mail: exp-pet@sceti.co.jp

Manufactured by

PPMX PERSEUS PROTEOMICS

Perseus Proteomics Inc.

4-7-6 Komaba, Meguro-ku, Tokyo 153-0041, Japan
URL: <http://www.ppmx.com>