

## Anti human SF-1 mouse monoclonal antibody

SF-1: Steroidogenic factor 1

Code No	PP-N1665-00
	old No. —
Clone No.	N1665
Lot.	A-2
Concentration	1 mg/mL
Volume	100 uL
Ig Class	G1

Description

Nomenclature NR5A1

Genbank

Specificity

Formulation

Origin

U76388

SF-1 (AD4BP, FTZF1,NR5A1) is still considered as an orphan nuclear receptor belongs to subfamily 5, It was found as a regulator of steroidogenic enzyme gene expression. Oxysterols are suggested as ligands. Its expressed in all sterodogenic tissues, including adrenal cortex, testicular Sertoli cells and Leydig cells, ovarian theca, hypothalamus and anterior pituitary. SF1 plays an important role in adrenal and gonadal development, such as hypothalamic-pituitary-gonadal axis and sex determination.

Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and

immunized with E.coli-expressed recombinant human

This antibody specifically recognizes human SF-1 and cross reacts with mouse and rat SF-1.

spleen cells derived from a BALB/c mouse


## Application / Recommended Concentration


In order to obtain the best results, optimal working dilutions should be determined by each individual user.

Western Blot	3 ug/mL
Non reducing Western Blot	3 ug/mL
ELISA	0.1 ug/mL (A450=1.0)
Immunoprecipitation	Decide by use
Supershift Assay	Not yet tested
Chromatin immunoprecipitation	Not yet tested

Immunohistochemistry

10 ug/mL





Rat		
Adrenal gland		
paraffin section		

Rat Salivary gland paraffin section

## Storage

Store at 2 - 8 °C up to one month. For long-term storage, the solution may be frozen in working aliquots. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended.

Reference

Jae Mi Suh, et al. Mol Endocrinol, Dec. 2006, 20(12): 3412 -3420

Jun Qin, et al. Developmental Dynamics, 2007, 236: 810-820

Purification Ammonium sulfate fractionation

SF-1(220-461aa)

Physiological saline with 0.1% NaN3 as a preservative.

**Notes** 

Sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large amounts of water during disposal.

## FOR RESEARCH ONLY. NOT FOR USE IN HUMANS.

Not for Diagnostic or Therapeutic use. Purchase of this product does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written consent of Perseus Proteomics Inc. is prohibited. MADE IN JAPAN Apr 24, 2008

Distributed by



3-6-7 Kasumigaseki, Chiyoda-ku, Tokyo 100-0013, Japan Tel +81(3) 5510-2347 Fax +81(3) 5510-0133 URL: http://www.sceti.co.jp/export/ e-mail: exp-pet@sceti.co.jp



Manufactured by